

## **Hysterectomies:A Retrospective study in Subharti Medical College, Meerut.**

**Mamta Tyagi<sup>1</sup>, Divya Mangla<sup>2</sup> and J.K. Goel<sup>3</sup>**

### **ABSTRACT**

Hysterectomy is the most common operation performed in Obstetrics and Gynaecology department. This is a retrospective study compiling various indications, demographic details and type of operation performed in 1065, cases of hysterectomy performed in the department of Obstetric and Gynaecology with in 4 years from 2005 – 2008 in Subharti Medical College & Hospital.

Analysis revealed that 1055 hysterectomies (99.06%) were performed for Gynaecological indications while the rest 10 (.94%) hysterectomies were performed for Obstetrical indications. Among Gynaecological hysterectomies, vaginal hysterectomies (632 ,(59.9%) were much more frequent than abdominal hysterectomies 423 (40.1%). Commonest Gynaecological indication was DUB (34.28%) and the main Obstetrical indication was rupture uterus (60.0%). Most common symptom for gynaecological indication was menstrual disturbances 675 (64.0%), and among menstrual disturbances, menorrhagia was commonest. Common age group was 40 – 49 years for gynaecological hysterectomies and 30 – 39 years for Obstetrical hysterectomies.

**Key Words :** Laproscopic hysterectomy, demographic, obstetrical